

Central catheter of peripherally insertion in neonates: integrative literature review

Rangel, Regiane Josy Mediate; Castro, Denise Silveira de; Primo, Cândida Caniçali; Zandonade, Eliana; Christoffel, Marialda Moreira; Amorim, Maria Helena Costa

Veröffentlichungsversion / Published Version
Zeitschriftenartikel / journal article

Empfohlene Zitierung / Suggested Citation:

Rangel, R. J. M., Castro, D. S. d., Primo, C. C., Zandonade, E., Christoffel, M. M., & Amorim, M. H. C. (2016). Central catheter of peripherally insertion in neonates: integrative literature review. *Revista de Pesquisa: Cuidado é Fundamental Online*, 8(4), 5193-5202. <https://doi.org/10.9789/2175-5361.2016.v8i4.5193-5202>

Nutzungsbedingungen:

Dieser Text wird unter einer CC BY-NC Lizenz (Namensnennung-Nicht-kommerziell) zur Verfügung gestellt. Nähere Auskünfte zu den CC-Lizenzen finden Sie hier:
<https://creativecommons.org/licenses/by-nc/4.0/deed.de>

Terms of use:

This document is made available under a CC BY-NC Licence (Attribution-NonCommercial). For more Information see:
<https://creativecommons.org/licenses/by-nc/4.0>

Cateter central de inserção periférica em neonato: revisão integrativa da literatura

Central catheter of peripherally insertion in neonates: integrative literature review

Catéter central de inserción periférica en el neonato: revision integrativa de la literatura

Regiane Josy Mediate Rangel¹, Denise Silveira de Castro², Cândida Caniçali Primo³, Eliana Zandonade⁴, Marialda Moreira Christoffel⁵, Maria Helena Costa Amorim⁶

How to quote this article:

Rangel RJM; Castro DS; Primo CC, et al. Central catheter of peripherally insertion in neonates: integrative literature review. Rev Fund Care Online. 2016 out/dez; 8(4):5193-5202. DOI: <http://dx.doi.org/10.9789/2175-5361.2016.v8i4.5193-5202>

ABSTRACT

Objective: To evaluate the evidence published on the practices of the use of Peripheral Inserted Central Venous Catheter (PICC) in newborns. **Method:** Integrative review researching the databases LILACS, MEDLINE, and COCHRANE, from 2008 to 2012. **Results:** Forty one articles were selected and ranked by evidence level: 4.9% level one; 9.8% level two; 31.7% level three; 34.2% level four; 17% level five; and 2.4% level six. These articles were grouped into the categories: general complications; blood flow infection related to catheter; utilization description and insertion of practices and maintenance. **Conclusion:** The evidence highlighted the importance of the permanent education for the insertion, maintenance and application of new technologies, in order to

¹ Enfermeira especialista em neonatologia, mestra em Enfermagem. Atua como enfermeira assistencial na Unidade de Terapia Intensiva Neonatal do Hospital Universitário Cassiano Antonio de Moraes, Vitória, ES.

² Possui graduação em Enfermagem pela Universidade Federal do Espírito Santo (1980), mestrado em Enfermagem pela Universidade de São Paulo (1990) e doutorado em Enfermagem pela Universidade Federal do Rio de Janeiro (1999). Atualmente é professor adjunto aposentada da Universidade Federal do Espírito Santo. Coordenadora Adjunta do Doutorado Interinstitucional UFES/UFRJ - DINTER e Professora Permanente do quadro de docentes do Programa de Pós Graduação em Enfermagem UFES.

³ Possui graduação em Enfermagem pela Universidade Federal do Espírito Santo (1997) e mestrado em Atenção à Saúde Coletiva pela Universidade Federal do Espírito Santo (2005). Atualmente é professor assistente da Universidade Federal do Espírito Santo.

⁴ Possui graduação em Matemática (1990) e Estatística (1994) pela Universidade Federal do Espírito Santo, mestrado em Teoria de Controle e Estatística pela Pontifícia Universidade Católica do Rio de Janeiro (1993) e doutorado em Estatística pela Universidade de São Paulo (1999). Atualmente é professora associada do Departamento de Estatística da Universidade Federal do Espírito Santo, bem como professora permanente do Programa de pós graduação em Saúde Coletiva na mesma universidade. E-mail: elianazandonade@uol.com.br

⁵ Possui graduação em Enfermagem pela Universidade Estadual do Ceará (1987), mestrado em Enfermagem pela Universidade Federal de São Paulo (1994) e doutorado em Enfermagem pela Universidade Federal do Rio de Janeiro (2002). Pós-doutorado Escola de Enfermagem Ribeirão Preto - EERP-USP. Centro Colaborador da Organização Mundial de Saúde para o desenvolvimento da Pesquisa em Enfermagem. Atualmente é professor adjunto da Universidade Federal do Rio de Janeiro.

⁶ Possui graduação em Enfermagem e Obstetrícia pela Universidade Federal do Espírito Santo (1983), especialização e mestrado na área de Enfermagem Obstétrica pela Universidade Federal de São Paulo (1987) e doutorado em Enfermagem pela Universidade Federal do Rio de Janeiro (1998). Atualmente é professor associado da Universidade Federal do Espírito Santo do Curso de Graduação em Enfermagem, Coordenadora do Programa de Pós-graduação em Enfermagem, docente do Programa de Pós-Graduação em Saúde Coletiva

minimize the unwanted effects of the use of PICC. It was noted that there is a lack of national production of studies that showed strong evidence levels.

Descriptors: Central Venous Catheterization, Newborn, Neonatal Intensive Care Unit, Neonatal Nursing.

RESUMO

Objetivo: Avaliar as evidências publicadas acerca das práticas no uso do Cateter Venoso Central de Inserção periférica (PICC) em Recém-Nascido. **Método:** Revisão integrativa que pesquisou as bases de dados LILACS, MEDLINE e COCHRANE nos anos de 2008 a 2012. **Resultados:** Selecionaram-se 41 artigos, os quais foram classificados, em nível de evidência, 4,9% nível um, 9,8% nível dois, 31,7% nível três, 34,2% nível quatro, 17% nível cinco, 2,4% nível seis. Foram agrupados nas categorias: complicações gerais, infecção de corrente sanguínea relacionada ao cateter, descrição da utilização e práticas de inserção e manutenção. **Conclusão:** As evidências ressaltaram a importância da educação permanente para inserção, manutenção e aplicação de novas tecnologias para minimizar os efeitos indesejados do uso do PICC. Observou-se escassez de produção nacional de estudos que retratem fortes níveis de evidências.

Descritores: Cateterismo venoso central, Recém-nascido, Unidades de Terapia Intensiva Neonatal, Enfermagem Neonatal.

RESÚMEN

Objetivo: Evaluar las evidencias publicadas acerca de las prácticas del uso del Catéter Venoso Central de Inserción Periférica (PICC) en Recién Nascidos. **Método:** Revisión integrativa que investigó las bases de datos LILACS, MEDLINE y COCHRANE en los años de 2008 a 2012. **Resultados:** Se seleccionaron 41 artículos, que fueran clasificados, en nivel de evidencia, 4.9% nivel uno, 9.8% nivel dos, 31.7% nivel tres, 34.2% nivel cuatro, 17% nivel cinco, y uno 2.4% nivel seis. Fueran agrupados en las categorías: complicaciones generales, infección de corriente sanguínea relacionada al catéter, descripción de la utilización y de las prácticas de inserción y mantenimiento. **Conclusión:** Las evidencias resaltarán la importancia de la educación permanente para la inserción, mantenimiento y aplicación de nuevas tecnologías para minimizar los efectos indeseados del uso del PICC. Fue observado la escasez de producción nacional de estudios que retraten fuertes niveles de evidencias.

Descriptores: Cateterismo venoso central, Recién Nacido, Unidades de Terapia Intensiva Neonatal, Enfermería Neonatal.

INTRODUCTION

Scientific and technological advances are increasing in the area of neonatology, as well as assistance in the Neonatal Intensive Care Unit (NICU), which contribute to a significant increase in the survival rate of newborn preterm and low-weight, changing the newborn mortality profile.¹

In nursing care of neonatology, they highlight the need for safe and durable venous access for administration of antibiotics, intravenous hydration, parenteral nutrition, vasoactive drugs and others.²⁻³

Currently an alternative widely used in a stable and effective venous access for newborns critically ill in NICUs is Central Catheter of Peripheral Insertion (PICC), so that the catheter is long and flexible, inserted through a peripheral vein, it progresses to the distal third of the cava vena – superior or inferior, thus gaining access for the central venous.⁴⁻⁷

However, the use of this device is associated with some complications that can occur during insertion, maintenance and removal, but often less frequent than other catheters of central location.⁸⁻¹⁰

Despite its benefits, the use of PICC is a complex process that requires technical and scientific knowledge specific to their insertion and maintenance, requiring skilled professionals and employing safe procedures and technologies to minimize side effects while using this device.¹¹⁻²

Professionals responsible for the appointment, insertion and maintenance of PICC must provide special care related to technical and technologies that promote success in catheter insertion, perform the appropriate management of pain control, use methods of visualization and optimal positioning of the catheter tip, take measures to avoid the main complications and prevention of infection related to catheter.¹³⁻⁵

To accompany these developments and encourage improvements in the quality of care provided to the differentiated clientele, it is important a reflective practice based on scientific knowledge, as evidence-based practice, which seeks to find in search results the main ideals and efficient conduct of a problem established through the organization of consistent and relevant evidence.¹⁶

Before the above, this review aims to assess the available evidence about the practices in the use of PICC in newborns and its implications, as the production of scientific knowledge about this practice can contribute to the development and incorporation of strategies for patient safety and quality of care to high-risk newborns that contribute to the reduction of neonatal mortality.

METHODS

This is an integrative review of the literature, considered as a research method that enables the search, critical evaluation and synthesis of the state of knowledge about a particular topic or issue in a systematic and orderly manner, and point to the implications in the professional practice and gaps in the scientific production that need to be filled with the realization of new studies.¹⁷

For the realization of this integrative review, the steps proposed by Ganong were as follows: (1) issue identification and selection guiding question; (2) literature search with the establishment of criteria for inclusion and exclusion of studies; (3) data collection with defining the information to be extracted from selected studies and categorization of studies; (4) Critical analysis of the included studies; (5)

interpretation and discussion of results and (6) presentation of the review/synthesis of knowledge.¹⁷

It was established the following question: what are the published evidence about the practices in the use of PICC in newborns and its implications?

A search of the scientific literature according to the following inclusion criteria was performed: productions in Portuguese, English or Spanish, included from January 2008 to December 2012, available in the databases Latin American and Caribbean Sciences Health (LILACS), Medical Literature Analysis and Retrieval System online (Medline) and Cochrane Library (Cochrane).

For electronic search, the following descriptors were used in Health Sciences (DeCS), "central venous catheterization," "peripheral catheterization" and "newborn".


The data were organized as item identification, institution of study, type of publication, features and methodological rigor.

To determine the level of evidence, the following hierarchy proposal of evidence was used:¹⁸

- Level 1: evidence from meta-analysis of multiple controlled and randomized clinical trials;
- Level 2: evidence from individual studies with experimental design;
- Level 3: evidence of quasi-experimental studies, such as studies without randomization with one group pre and post-test, time series or case control;
- Level 4: evidence from descriptive studies (non-experimental) or qualitative approach;
- Level 5: evidence from case reports or experience;
- Level 6: evidence based on expert opinions.

The figure 1 shows the flowchart of the sample selection of the studies included in the integrative review.

Figure 1 - Flowchart of the sample selection of the studies included in the Integrative Review - March/2013 (RANGEL, CASTRO, PRIMO, 2013)



RESULTS AND DISCUSSION

They found 470 articles (117 LILACS 191 MEDLINE and COCHRANE 172), being excluded 424 which did not meet the inclusion criteria and included 46 articles. From these, 5 had up in duplicate, for a total of 41 final sample articles.

After exhaustive reading of the 41 items selected in the final sample, we elaborated the themes: insertion practices and maintenance of PICC (18 articles), description of the use of PICC (07 articles), general complications of PICC (09 articles) and bloodstream infection related to catheter (07 articles).

We conducted a synthesis of articles about the category, authorship, journal, country, language, year and level of evidence, presented in Figure 2.

Among the articles selected, 29.3% were published in Brazilian journals in Portuguese, 24.4% in US journals in English, 46.3% in international journals of different countries, as Turkey, India, England, Taiwan, Japan, the Netherlands, and all in English. There was 20% publications in 2012; 22.5% in 2011; 27.3% in 2010; 7.7% in 2009; and 22.5% in 2008.

As for the level of evidence, 4.9% had level one, level two 9.8%, 31.7% level three, level four 34.2%, 17% level five, and 2.4% level six. The highlight was the strongest level of evidence (1, 2 and 3) in international studies, still incipient in national, which were predominantly descriptive.

Considering the importance of strong evidence for the practice, there was the synthesis of evidence items 1, 2 and 3, which are shown in Figures 3, 4 and 5, respectively, as written, study type, action, outcomes, and conclusions.

Figure 2 - Summary of articles about the category (Cat), authorship, journal, country, language, year and level of evidence (LE). USA: The United States of America. Vitória 2013 (RANGEL, CASTRO, PRIMO, AMORIM, CHRISTOFFEL, ZANDONADE, 2013)

Cat	Author	Journal	Country	Country	Year	LE
PRACTICES OF INSERTION AND MAINTENANCE	Sartoli et al ¹⁹	Nursing (São Paulo)	Brazil	Brazil	2012	4
	Jain et al ²⁰	Am J Perinatol	USA	USA	2012	3
	Johann et al ²¹	Rev Esc Enferm USP	Brazil	Brazil	2012	5
	Marcatto et al ²²	Arch Dis Child Fetal Neonatal	Brazil	Brazil	2011	2
	Taylor et al ²³	Adv Neonatal Care	USA	USA	2011	3
	Fidler et al ²⁴	Adv Neonatal Care	USA	USA	2011	1
	Uygun et al ²⁵	Acta Cir. Bras	Turkey	Turkey	2011	3
	Uslu et al ²⁶	J Perinatol	Turkey	Turkey	2010	2
	Johann et al ²⁷	REME rev. min.	Brazil	Brazil	2010	4
	Sharpe et al ²⁸	Adv Neonatal Care	USA	USA	2010	5
	Corzine et al ²⁹	Neonatal Netw	USA	USA	2010	5
	Monsky et al ³⁰	J Vasc Access	USA	USA	2010	4
	Sneath et al ³¹	Neonatal Netw	USA	USA	2010	5
	Ragavan et al ⁷	Indian J Pediatr	India	India	2010	3
	Smirk et al ³²	Arch Dis Child Fetal Neonatal	Australia	Australia	2009	2
	Lago et al ³³	Paediatr Anaesth	Italy	Italy	2008	2
	Shah et al ³⁴	Cochrane Database Syst Rev	Canada	Canada	2008	1
	Chaves et al ³⁵	Nursing (São Paulo);	Brazil	Brazil	2008	6
GENERAL COMPLICATIONS	Costa et al ¹²	Rev. Gaúcha Enferm	Brazil	Brazil	2012	4
	Kalkra et al ³⁶	J Vasc Access	England	England	2012	5
	Tosello et al ³⁷	J Pediatr Surg	France	France	2011	5
	Tsai et al ³⁸	Infect Control Hosp Epidemiol	Taiwan	Taiwan	2011	3
	Bulbul et al ³⁹	J Matern Fetal Neonatal	Turkey	Turkey	2010	4
	Wolfe et al ⁴⁰	Adv Neonatal Care	USA	USA	2010	5
	Francheschi et al ¹¹	Rev Lat Am Enfermagem	Brazil	Brazil	2010	4
	Liu et al ⁴¹	J Infus Nurs	China	China	2009	3
	Ohki et al ⁵	Pediatr Int	Japan	Japan	2008	3

(To be continued...)

(Continuation)

Cat	Author	Journal	Country	Country	Year	LE
INFECTION	Ponnusamy ⁴²	Arch Dis Child Fetal	England	England	2012	3
	Tsai et al ⁴³	Am J Infect Control.	Taiwan	Taiwan	2012	3
	Njere et al ⁴⁴	J Pediatr Surg	England	England	2011	3
	Hsu et al ⁴⁵	Pediatr Neonatol	Taiwan	Taiwan	2010	3
	Van den Hoogen et al ⁴⁶	Acta Paediatr	Netherlands	Netherlands	2008	3
	Garland et al ⁴⁷	Infect Control Hosp Epidemiol	USA	USA	2008	3
	Hoang et al ⁴⁸	Pediatrics	USA	USA	2008	4
DESCRIPTION OF USE	Ishida et al ⁴⁹	Nursing (São Paulo)	Brazil	Brazil	2012	4
	Dórea et al ⁵⁰	Rev Bras Enferm	Brazil	Brazil	2011	4
	Reis et al ¹⁰	Rev Bras Enferm	Brazil	Brazil	2011	4
	Baggio et al ¹⁴	Rev Gaucha Enferm	Brazil	Brazil	2009	4
	Freitas et al ⁵¹	REME rev. min. enferm	Brazil	Brazil	2009	4
	Camargo et al ⁶	Rev Esc Enferm USP	Brazil	Brazil	2008	4
	Bueno et al ⁴	J Perinatol	Spain	Spain	2008	4

Figura 3 - Summary of evidence of articles Level 1 as the authorship, study type, intervention, results and conclusions. PICC: Central Catheter of Peripherally Insertion; NB: Newborn (RANGEL, CASTRO, PRIMO, AMORIM, CHRISTOFFEL, ZANDONADE, 2013)

Author/Type of study	Intervention	Results and conclusion
Fidler <i>et al.</i> ²⁴ Meta-analysis	Use of bedside ultrasound during insertion and placement of the PICC.	It has been associated with: increased success rate on the first try; declines in thrombosis rates; increased number of options veins; ability to put large-caliber catheter; ability to view and prevent arterial puncture; decreased tissue trauma; affordable equipment; experienced professional requirement
Shah <i>et al.</i> ³⁴ Meta-analysis	Use of prophylactic heparin.	It reduces occlusion and enables greater number of patients completing the therapy desired. It supports the prophylactic use of heparin for PICC in newborns, at a dose of 0.5 IU/kg/h. None of these studies was designed to evaluate the rate of adverse events.

Figura 4 - Summary of evidence of articles Level 2 as the authorship, study type, intervention, results and conclusions. EMLA: eutectic mixture of lidocaine and prilocaine; PICC: Central Catheter Peripherally Insertion; NB: Newborn (RANGEL, CASTRO, PRIMO, AMORIM, CHRISTOFFEL, ZANDONADE, 2013)

Author/Type of study	Intervention	Results and conclusions
Marcatto <i>et al.</i> ²² Randomized clinical trial	Use of oral glucose 25% or EMLA cream in pain management in preterm infants undergoing insertion of PICC.	There were no statistically significant differences between the groups for the primary outcome. Other strategies should be considered to control pain during this procedure.
Uslu <i>et al.</i> ²⁶ Clinical, prospective, randomized, controlled, double-blind	Use of heparin in a low dose on permeability and occlusion of the PICC in newborns.	In the group of heparin continuous infusion of low dose heparin (0,5 IU kg ⁻¹ h ⁻¹) the duration of catheter patency was higher and catheter occlusion rate was smaller and allowed the completion of therapy without increasing adverse effects.
Smirk <i>et al.</i> ³² 2009 Randomized clinical trial	Pressure changes in neonatal PICC under different laboratory conditions and the risk of rupture in the clinical practice.	The polyurethane PICC has tolerance of pressure higher than the silicone catheters and is less likely to break under experimental conditions. Silicone Catheters clogged easily break when they receive flush.
Lago <i>et al.</i> ³⁵ Randomized clinical trial	Infusion of low dose of remifentanyl during PICC in premature infants.	Validated pain scales, cardiovascular movements, respiratory response and body during insertion of PICC suggest control of pain and discomfort with remifentanyl, but the time to complete the insertion and the number of attempts required remained the same.

Figura 5 - Summary of evidence of articles Level 3 as the authorship, study type, intervention, results and conclusions. CRBSI: bloodstream infection related to catheter; PICC: Central Catheter

Author/Type of study	Intervention	Results and conclusions
Ponnusamy <i>et al.</i> ⁴² Observational	Culture of the middle and proximal segments of the catheter beyond the tip.	The culture of the middle, proximal and tip segments did not improve the diagnosis of CRBSI in relation to only tip culture.
Jain <i>et al.</i> ²⁰ Comparative	Using x-rays to determine PICC tip position compared with the use of echocardiography in the NICU.	Echocardiography is useful tool for tip position identification, making manipulation in real time, and minimizing the exposure of new X-rays.
Tsai <i>et al.</i> ⁴³ Retrospective cohort study	It assesses removal of the PICC in newborns with CRBSI, and examines the risk factor for infectious complications.	The retention of the catheter for more than 3 days in NB with CRBSI is associated with a delay resolution and a higher incidence of recurrence within one month.
Njere <i>et al.</i> ⁴⁴ Retrospective cohort study	It identifies the rates of complications and possible predictors of infection related to the PICC in a NICU	The PICC infection rate was 17/1000 catheter-days. OR = 3,1 catheter in situ for 9 days or more. Negative coagulase staphylococcus was isolated in 89% blood cultures.
Taylor <i>et al.</i> ²³ Retrospective control case	Establishment of a team dedicated to PICC and CRBSI.	The CRBSI was reduced by almost half after the creation of a team dedicated to PICC.
Tsai <i>et al.</i> ³⁸ Retrospective cohort study	Insertion time and PICC permanence over the complications.	Insertion with more than 60 minutes and length of stay for more than 30 days is associated with higher rates of complications related to the catheter.
Uygun <i>et al.</i> ²⁵ Retrospective cohort study	Insertion time and PICC permanence over the complications.	95% of the PICC have been successfully inserted. Major complications were observed. The new insertion technique can be a safe and easy procedure.
Liu <i>et al.</i> ⁴⁰ Retrospective, comparative descriptive	It analyzes complication rates with PICC in a NICU and study the risk factors associated with non-elective catheter removal.	The complication rate was 31.73%, with most of phlebitis. The NB of elective removal group had position in the vena cava. Mechanical complications are the most common reasons for removal of catheters.
Ohki <i>et al.</i> ⁵ National questionnaire to NICU in Japan	Institutional policies regarding the use of PICC and also the frequency of complications.	Tip positions of the catheter out of the heart were preferred. The frequency of pericardial effusion and cardiac tamponade were 0.07 to 0.11%.
Garland <i>et al.</i> ⁴⁷ Retrospective cohort study	It defines the pathogenesis of CRBSI in NB with PICC.	Most CRBSI was caused by coagulase-negative staphylococci and derive from intraluminal contamination (67%).
Hsu <i>et al.</i> ⁴⁵ Descriptivo comparativo	It assesses the risk of infection and other complications associated with PICC in newborns with very low weight.	Significant risk factors of CRBSI include catheters inserted in the femoral sites and a longer duration of placement of PICC.
Ragavan <i>et al.</i> ⁷ Comparative descriptive	It compares complication rates of surgical central venous catheters (CL) and PICC.	The rates of all complications were significantly lower in the PICC group. The removal before the completion of therapy was in 11.5% (PICC) and 37.5% (CL).
Van den Hoogen <i>et al.</i> ⁴⁶ Comparative descriptive	It assesses effect of antibiotic administration at the time of removal of the PICC	The administration of antibiotics at the time of catheter removal reduced significantly the incidence of sepsis.

Insertion And Maintenance Practices Category

The articles in this category discuss about technology, techniques and care used in the practice of inserting and maintaining the PICC.

The insertion of the catheter in newborns is a very delicate procedure, because of the fragility of this patient, small veins and need thin catheters.⁴⁶

In studies with purposes of analyzing new insertion techniques they presented in the insertion a success rate of 95%, without observing significant complications, demonstrating that are safe and easy procedure that can be used as an alternative for insertion of the PICC.^{4,46}

Also as a facilitator for successful integration, the use of ultrasound (US) of the head during insertion and placement of the PICC was described in meta-analysis associated with: increased success rate in the first attempt; increased number of options veins; ability to put large-caliber catheter; ability to view and prevent arterial puncture; decreased tissue trauma; and more affordable equipment. However, this requires experienced professional with training and eye-hand coordination.²⁴

Regarding pain control in insertion procedure, the remifentanyl in low dose has measurable and synergistic analgesic effect in combination with 12% sucrose and non-nutritive sucking, but it makes the insertion of PICC be easier or more fast.³³ The suction with glucose at 25% and EMLA cream alone were not sufficient strategies to control pain.²²

The position of the tip of the catheter to its proper position is essential to prevent complications. This should be located in the superior vena cava or inferior vena cava near the junction with the right atrium, 0.5 to 1 cm out of the heart chamber of NB.^{20,28,31}

It is not clear what the best viewing method of literature review localization.^{20,31} A research states that fluoroscopy is ideal, but cannot be performed at the bedside and has a high cost, and the supine chest radiograph is more convenient and more used.³¹

However, a comparative study concluded that the use of echocardiography has increased accuracy and minimizes exposure to x-rays, and also claims that the arm position is very important in performing radiography because their movement can cause catheter migration.²⁰ The movements of arm can even be used in non-invasive techniques to reposition the tip.²⁸

In relation to care for maintaining the catheter they emphasized the use of heparin to prevent obstruction, aseptic dressing technique and the pressure applied to the catheter and prevent disruption.^{19,21,26,29,34}

The continuous infusion of low dose of heparin (0.5 IU/kg/h) into the fluids is an effective measure to reduce the occlusion of the catheter, allowing the completion of therapy without observing adverse effects.^{26,34}

The healing practice is essential in maintaining the PICC because it covers and prevents local trauma and contamination, and respect the aseptic technique and

the evidence as to the material used and the frequency of change.²¹ In reporting experience with 491 catheters that evaluates technique of dressing with a protective base layer they concluded that catheter complication rates were low, and that dressing changes catheters wereminimized.³⁷

As for the pressure applied to the catheter and the risk of rupture, only one study addressed the issue.³² A randomized clinical trial examined the pressure changes in neonatal PICC in different laboratory conditions and found that the polyurethane have tolerance of pressure greater than the silicone catheters and they are less likely to break under experimental conditions. The silicone catheters clogged are broken easily when they receive flush even with syringe in a greater caliber.³²

Faced with various evidence and conduct the insertion and maintenance of the PICC, the standardization of everyday practice in the care of newborns is critical. An experience report presented a form to unify the practices related to the PICC in a NICU, which has been a supporting tool for early detection of alterations that result in risk of complications for newborns, contributing to a qualified care.⁴²

In addition to standardizing the behavior, the establishment of a team dedicated to PICC also improves the quality of care, reaching to reduce by almost half the CRBSI in infants with extremely low birth weight requiring venous access in a long term.²⁹

In conclusion, this category emphasizes the importance of skills training for insertion and maintenance as well as the need to implement new technologies to minimize the undesirable effects in the use of PICC.

General surgery category

it highlights studies that provide information about the complications associated with the use of PICC in newborns.

Prevalence of complications has been described in different studies, ranging from 30.7% to 50.7% (30,7%⁴⁸, 31,7%⁴⁰, 39,3%¹², 45,9%³⁹, 47,6%¹¹, 50,7%³⁸). Despite the differences between these values and the reality of each search location, it can be seen that the prevalence show an average of 41.1% of complications.

The most common complications were reported in studies:

- Obstruction (6,9%⁴⁸; 12,7%³⁹; 13,1%¹²; 16,6%³⁸; 19,4%¹¹);
- Disruption (7,1%³⁸; 8,8%¹¹; 9,5%¹²; 11,2%²⁶; 15,4%⁵⁰)
- Phlebitis (5,6%⁴⁸; 10,8 %³⁸; 22,1%⁴⁰)
- Limb edema (7,1%¹²)
- Infection related to PICC (2,3%¹¹; 6%¹²; 11,1 %⁴⁸; 36,4%³⁸)
- Extravasation (1,2%¹²; 4,3%³⁸; 5,8%⁴⁸);
- Accidental Attraction (1,2%¹²; 2,3%¹¹)

Regarding the outcome, complications often lead to non-elective removal of PICC.^{12,40} On the other hand, they have also been reported some rare complications such as

diaphragmatic paralysis, oliguria with the presumed etiology of tip catheter misplaced which blocks renal vein and hypertonic solution and parenteral nutrition administration directly into the renal vein, the disruption and migration of the catheter to the pulmonary artery, pericardial effusion, cardiac tamponade, pleural effusion, ascites and difficulty of removal.^{5,12,29,37,39,41}

Some authors concluded that most of the complications were related to care, requiring qualification of professionals to better care related to insertion technique, maintenance and management of catheter.^{11-2,36}

It also stands out in this category more evidence from international research indicating the need for national studies that address the theme and have a strong level of evidence.

Whereas the infection was a major complication of the use of PICC and showed up seven articles dealing exclusively with this question, so we constituted a new topic category: infection.

Infection category

the infection category contains seven articles that have as main theme the bloodstream infection related to catheter (CRBSI) in newborns.

Regarding the definition of CRBSI studies we used similar concepts, being defined as: positive culture of at least one blood sample obtained from a peripheral vein, clinical signs of infection, no other site of infection and PICC insertion at least 5 days.⁴²⁻⁷

The CRBSI can be confirmed through the tip culture catheter. Thus, prospective observational study in England with 189 catheters in 143 newborns, 47 with suspected CRBSI, concluded that the culture of the middle segments, proximal and the tip catheter has not improved the diagnosis of infection in relation to only tip culture.⁴²

The principal pathogen related to CRBSI found in blood cultures, this was the negative-coagulase staphylococci (CNS), as noted in a prospective cohort study of 218 newborns and 294 catheters, conducted in England, in which isolated the ECN from the 89% of 62 blood cultures, followed by *Klebsiella* (3.2%), *Escherichia coli*, *Candida*, *Enterococcus* and *Staphylococcus aureus* (1.6% each).⁴⁴

Similarly, another study had ECN isolated in 40.1% of 67 positive blood cultures, followed by *Staphylococcus aureus* (16.4%), *Enterococcus* and *Klebsiella* (13.4%).⁴⁵ Another study also showed the presence of ECN in 78% of the positive blood cultures and 92% had a positive culture of the catheter segments that corroborate with similar results of other authors.^{42,46-7}

Regarding the PICC infection rate, there are two articles, one obtained rate of 8.3 per 1000 catheter days, while the other obtained rate of 17 per 1000 catheter in a day.⁴³⁻⁴

In relation to the insertion site, a study found in a sample of 396 infants and 477 catheters (370 upper limb and 107 lower) a CRBSI rate of 11.6% for the superior and 9.3% in

catheters in inferior limbs.⁵ Another survey, with a sample of 292 infants and 412 catheters, found rates of 12.4% for non-femoral site and 21.6% for femoral site (increased risk of CRBSI compared with non-femoral catheters of 1,76).⁴⁵ More studies are needed to clarify the relationship of the insertion site with CRBSI in newborns.

Besides the insertion site, another predictor of infection that stood out was the duration time of catheter. Research has found an odds ratio of 3.1 and $P < 0.01$ for an infection by PICC if the catheter was in situ for 9 days or more.⁴⁴

Regarding the pathogenesis of CRBSI of PICC, only one study evaluated the derivation of contamination and concluded that most have intraluminal contamination source. Prospective cohort study nested with 82 neonates identified 15 CRBSI, 67% acquired through intra luminal, 20% extra luminal and 13% undetermined. To this predominant mechanism of infection, strategies for prevention are more likely to be effective.⁴⁷

With regard to indication to remove or not the catheter of RN who requires venous access in term of diagnosis of CRBSI, we concluded that the PICC should be removed, because the retention of PICC for more than three days is associated with delayed of resolution of clinical sepsis and increased incidence of recurrence within a month.⁴³

Also regarding the removal, it was observed that the administration of antibiotics for the PICC removal, there were significant results in reducing incidence of sepsis after removal of catheter;⁴⁶ however prospective studies are needed to confirm this observation.

Despite various evidence presented in this category, the study highlights the need for further studies to confirm the existing results and clarify the questions and controversies still unanswered.

For this category we observed national studies. Thus, the scientific production on this theme is a challenge to Brazilian researchers.

Description picc use category

this category includes articles describing the use of the PICC, providing knowledge on the device's use of reality and the people who received it.

The characterization of population studies show that the majority are preterm infants, with low birth weight, male, appropriate for gestational age, primary diagnosis of respiratory disease, especially respiratory distress syndrome (RDS) due to prematurity.^{14, 49-51}

The most used type of silicone catheter was mono lumen.^{10,14,49,51} Most punctured venous catheters were located in upper limbs, especially the basilic vein, and then the cephalic.^{4,6,10,14, 49}

The positioning of the catheter tip, visualized by radiography, predominated in central location, especially in the superior vena cava.^{10,14,49,51} With respect to the PICC

residence time, the average of studies varied between 7.7 to 14, 5 days (7.7⁴⁹, 9.4⁵⁰, 10.5⁵¹, 13.5⁴, 14.5²⁹).

The predominant removal reason was the completion of the proposed therapy for the indication of PICC, followed by events and complications.^{4,6,14,47,49}

It was concluded that the evidence found on the characteristics of the use of PICC presents homogeneity despite being studies of different institutions and realities.

CONCLUSIONS

This integrative review of the published evidence about the use of practices of PICC in infants involved 41 articles which have the following levels of evidence: 4.9% level one, level two 9.8%, 31.7% level three, 34.2% level four, 17% level five, and 2.4% level six. The considered strong evidence (level 1, 2 and 3) amounted to 46.3%.

There was national production shortage of studies that portray strong levels of evidence with experimental design, considered the “gold standard” in evidence-based practice.

Most publications analyzed are international, available in English, which is a challenge to be overcome by the professionals involved with care, so the understanding and application of research results to understand the English language becomes crucial.

Other research on the use of PICC in newborns are needed to help with the decision-making to disputes as the best end of the display method, the use of antibiotics at the time of catheter removal; and support the implementation of new technologies, such as ultrasound and its practical applicability, contributing to patient safety and quality of care.

REFERENCES

1. Silva GRG, Nogueira MFH. Terapia intravenosa em recém-nascidos: orientações para o cuidado de enfermagem. Rio de Janeiro: Cultura Médica; 2004. p.23-37.
2. Infusion Nurses Society (INS) Brasil. Diretrizes práticas para a terapia intravenosa. São Paulo; 2008.
3. Ferreira FLC, Silva GF, Fonseca PML, Christoffel MM. Terapia intravenosa em neonatologia e na pediatria: uma revisão sistemática da literatura. Rev Cuidado é Fundamental, v. 2, p. 125-129, 2010.
4. Bueno TM, Diz AI, Cervera PQ, Pérez-Rodríguez J, Quero J. Peripheral insertion of double-lumen central venous catheter using the Seldinger technique in newborns. J Perinatol; 2008 Apr 28(4):282-.
5. Ohki Y, Yoshizawa Y, Watanabe M, Kuwashima M, Morikawa A. Complications of percutaneously inserted central venous catheters in Japanese neonates. Pediatr Int; 2008 Oct 50(5):636-9.
6. Camargo PP, Kimura AF, Toma E, Tsunehiro MA. Localização inicial da ponta de cateter central de inserção periférica (PICC) em recém-nascidos. Rev esc enferm USP; 2008; 42(4): 723-8.
7. Ragavan M, Gazula S, Yadav DK, Agarwala S, Srinivas M, Bajpai M, Bhatnagar V, Gupta DK. Peripherally inserted central venous lines versus central lines in surgical newborns--a comparison. Indian J Pediatr; 2010 Feb 77(2):171-4.
8. CDC. Guidelines for Environmental Infection Control in Health-Care Facilities. MMWR 2003; 52 (Nº. RR -10). Acessível em: http://www.cdc.gov/ncidod/dhqp/pdf/guidelines/Enviro_guide_03.pdf
9. Tariq, M; Huang, D. PICCing the best access for your patient. Crit Care, 2006 10(5).
10. Reis AT, Santos SBI, Barreto JM, Silva GRGd. O uso do cateter epicutâneo na clientela neonatal de um hospital público estadual: estudo retrospectivo. Rev. enferm. UERJ; 2011 out-dez 19(4): 592-597.
11. Franceschi ALT, Cunha MLCD. Adverse Events Related to the Use of Central Venous Catheters in Hospitalized Newborns. Rev. Latino-Am. Enfermagem, Ribeirão Preto, 2010 abr; 18(2):196-202. Disponível em: <http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0104-11692010000200009&lng=en&nrm=iso>. Acesso em 22 fev 2013. <http://dx.doi.org/10.1590/S0104-11692010000200009>.
12. Costa P, Kimura A F, Vizzotto MPS, Castro TED, West A, Dorea E. Prevalência e motivos de remoção não eletiva do cateter central de inserção periférica em neonatos. Rev. Gaúcha Enferm., Porto Alegre, 2012 set; 33(3). Disponível em: <http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1983-14472012000300017&lng=en&nrm=iso>. Acesso em 22 fev 2013. <http://dx.doi.org/10.1590/S1983-14472012000300017>.
13. Lourenço AS, Kakehashi TY. Avaliação da implantação do cateter central de inserção periférica em neonatologia. Acta Paul Enferm. 2003; 16(2):26-32.
14. Baggio MA, Bazzi FCS, Bilíbio CAC. Cateter central de inserção periférica: descrição da utilização em UTI Neonatal e Pediátrica. Rev. Gaúcha Enferm., Porto Alegre (RS) 2010 mar; 31(1):70-6. Disponível em: <http://seer.ufrgs.br/index.php/RevistaGauchadeEnfermagem/article/view/11693/8490> Acesso em 22 fev 2013.
15. Belo MPM, Silva RAM de C, Nogueira ILM, Mizoguti DP, Ventura CMU. Conhecimento de enfermeiros de Neonatologia acerca do Cateter Venoso Central de Inserção Periférica. Rev. bras. enferm., Brasília, v. 65, n. 1, Feb. 2012. Disponível em: <http://www.scielo.br/scielo.php?script=sci_arttext&pid=S003471672012000100006&lng=en&nrm=iso>. Acesso em 22 fev 2013. <http://dx.doi.org/10.1590/S0034-71672012000100006>.
16. Domenico EBL, Ide CAC. Evidence based nursing: principles and applicability. Rev Latino-am Enfermagem 2003 janeiro-fevereiro;11(1):115-8.
17. Ganong LH. Integrative reviews of nursing research. Research in Nursing & Health. 1987 Fev, 10(1), p 1-11.
18. Stetler CB, et al. Utilization-focused integrative reviews in a nursing service. Appl Nurs Res. 1998; 11(4):195-206.
19. Sartoli NR, Tessuto; M C, Almeida CBPDe. Manutenção e cuidados pós-inserção do PICC: revisão integrativa de literatura / Maintenance and care after PICC insertion: an integrative review of the literature. Nursing (São Paulo); 2012 out 15(173): 148-535.
20. Jain A, Mcnamara PJ, Ng E, El-Khuffash A. The use of targeted neonatal echocardiography to confirm placement of peripherally inserted central catheters in neonates. Am J Perinatol; 2012 Feb 29(2):101-6.
21. Johann DA, Lazzari LSMD, Pedrolo E, Mingorance P, Almeida TQRD, Danski MTR. Cuidados com cateter central de inserção periférica no neonato: revisão integrativa da literatura. Rev. esc. enferm. USP, São Paulo, 2012 dez 46(6). Disponível em: <http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0080-62342012000600030&lng=en&nrm=iso>. Acesso em 22 jun 2013. <http://dx.doi.org/10.1590/S0080-62342012000600030>.
22. Marcatto JDO, Vasconcelos PC, Araújo CM, Tavares EC, Pereira ESY. EMLA versus glucose for PICC insertion: a randomised triple-masked controlled study.. Arch Dis Child Fetal Neonatal, 2011 Nov 96(6).
23. Taylor T, Massaro A, Williams L, Doering J, McCarter R, He J, Talley L, Short B. Effect of a dedicated percutaneously inserted central catheter team on neonatal catheter-related bloodstream infection. Adv Neonatal Care; 2011 Apr 11(2):122-8.
24. Fidler HL. The use of bedside ultrasonography for PICC placement and insertion. Adv Neonatal Care; 2011 Feb 11(1):52-3.
25. Uygun I, Okur MH, Otcu S, Ozturk H. Peripherally inserted central catheters in the neonatal period. Acta Cir. Bras., São Paulo, 2011 Oct 26(5). Disponível em: <http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0102-86502011000500014&lng=en&nrm=iso>. Acesso em 22 fev 2013. <http://dx.doi.org/10.1590/S0102-86502011000500014>.
26. Uslu S, Ozdemir H, Comert S, Bolat F, Nuhoglu A. The effect of low-dose heparin on maintaining peripherally inserted percutaneous central venous catheters in neonates. J Perinatol; 2010 Dec 30(12):794-9.

27. Johann DA, Danski MTR, Pedrolo E, De Lazzari LSM, Mingorance P. Avaliação de um cuidado de enfermagem: o curativo de catéter central de inserção periférica no recém-nascido. *REME rev. min. enferm*; 2010 out-dez 14(4): 515-520.
28. Sharpe EL. Repositioning techniques for malpositioned neonatal peripherally inserted
29. Corzine M, Willett LD. Neonatal PICC: one unit's six-year experience with limiting catheter complications. *Neonatal Netw*; 2010 May-Jun 29(3):161-73.
30. Monsky WL, Garza AS, Loh S, Link DP. Snare technique for the placement of a peripherally inserted central catheter in the neonatal and pediatric patient. *J Vasc Access*; 2010 Apr-Jun 11(2):100-5.
31. Sneath N. Are supine chest and abdominal radiographs the best way to confirm PICC placement in neonates? *Neonatal Netw*; 2010 Jan-Feb 29(1):23-35.
32. Smirk C, Soosay Raj T, Smith AI, Morris S. Neonatal percutaneous central venous lines: fit to burst. *Arch Dis Child Fetal Neonatal Ed*; 2009 Jul 94(4):F298-300.
33. Lago P, Tiozzo C, Boccuzzo G, Allegro A, Zacchello F. Remifentanyl for percutaneous intravenous central catheter placement in preterm infant: a randomized controlled trial. *Paediatr Anaesth*; 2008 Aug 18(8):736-44.
34. Shah PS, Shah VS. Continuous heparin infusion to prevent thrombosis and catheter occlusion in neonates with peripherally placed percutaneous central venous catheters. *Cochrane Database Syst Rev*; 2008 (2):CD002772.
35. Chaves EMC, Câmara SMC, Araújo KRX, Feitosa TLO, Bezerra FSDM, Queiroz MVO. Cateter central de inserção periférica: ficha para recém-nascidos / Peripheral insertion of central venous catheter (PICC): a protocol to newborns. *Nursing (São Paulo)*; maio 2008 11(120): 230-234.
36. Kalra VK, Arora P, Lua J. Spontaneous fracture and migration of distal segment of a peripherally inserted central venous catheter to heart in a neonate. *J Vasc Access*; 2012 Jul-Sep; 13(3):403. <http://www.sciencedirect.com/science/article/pii/S1552885512000645>
37. Tosello B, Michel F, Merrot T, Chaumoitre K, Hassid S, Lagier P, Martin C. Hemidiaphragmatic paralysis in preterm neonates: a rare complication of peripherally inserted central catheter extravasation. *J Pediatr Surg*; 2011 Jul 46(7):E17-21.
38. Tsai M H, Chu S M, Lien R, Huang HR, Wang JW, Chiang Cc, Hsu JF, Huang YC. Complications associated with 2 different types of percutaneously inserted central venous catheters in very low birth weight infants. *Infect Control Hosp Epidemiol*; 2011 Mar 32(3):258-66.
39. Bulbul A, Okan F, Nuhoglu A. Percutaneously inserted central catheters in the newborns: a center's experience in Turkey. *J Matern Fetal Neonatal Med*; 2010 Jun 23(6):529-35.
40. Wolfe D M. A previously undescribed etiology for oliguria in a premature infant with a peripherally inserted central catheter. *Adv Neonatal Care*; 2010 Apr 10(2):56-9.
41. Liu H; Han T; Zheng Y; Tong X; Piao M; Zhang H. Analysis of complication rates and reasons for nonelective removal of PICCs in neonatal intensive care unit preterm infants. *J Infus Nurs*; 2009 Nov-Dec 32(6):336-40.
42. Ponnusamy V, Venkatesh V, Curley A, Musonda P, Brown N, Tremlett C, Clarke P. Segmental percutaneous central venous line cultures for diagnosis of catheter-related sepsis. *Arch Dis Child Fetal Neonatal Ed*; 2012 Jul 97(4):F273-8.
43. Tsai M H, Hsu J F, Lien R, Huang H R, Chiang C C, Chu S M, Liang H F, Huang Y C. Catheter management in neonates with bloodstream infection and a percutaneously inserted central venous catheter in situ: removal or not? *Am J Infect Control*; 2012 Feb 40(1):59-64.
44. Njere I, Islam S, Parish D, Kuna J, Keshtgar A S. Outcome of peripherally inserted central venous catheters in surgical and medical neonates. *J Pediatr Surg*; 2011 May 46(5):946-50.
45. Hsu JF, Tsai MH, Huang HR, Lien R, Chu SM, Huang CB. Risk Factors of Catheter-related Bloodstream Infection With Percutaneously Inserted Central Venous Catheters in Very Low Birth Weight Infants. *Pediatr Neonatol* 2010; 51(6):336-342. Disponível em: <http://download.journals.elsevierhealth.com/pdfs/journals/18759572/PIIS1875957210600653.pdf>. Acesso em 22 fev 2013.
46. Van Den Hoogen A, Brouwer MJ, Gerards LJ, Fler A, Krediet TG. Removal of percutaneously inserted central venous catheters in neonates is associated with the occurrence of sepsis. *Acta Paediatr*; 2008 Sep 97(9):1250-2.
47. Garland J S, Alex Cp, Sevallius JM, Murphy Dm, Good MJ, Volberding AM, Hofer L L, Maki DG. Cohort study of the pathogenesis and molecular epidemiology of catheter-related bloodstream infection in neonates with peripherally inserted central venous catheters. *Infect Control Hosp Epidemiol*; 2008 Mar 29(3):243-9.
48. Hoang VSJ, Chandler M, Busaloni E, Clifton-Koeppel R, Modanlou H D. Percutaneously inserted central catheter for total parenteral nutrition in neonates: complications rates related to upper versus lower extremity insertion. *Pediatrics*. 2008;121(5):e1152-9
49. Ishida ACH, Teixeira GT, Boaventura SGDS, Gabas VP. Avaliação da implantação do cateter venoso central de inserção periférica em neonatos / Evaluation of the implementation of the peripherally inserted central catheter in neonates. *Nursing (São Paulo)* 2012 mar;14(166): 152-157.
50. Dórea E, De Castro TE, Costa P, Kimura A F, Dos Santos FM. Práticas de manejo do Cateter Central de Inserção Periférica em uma unidade neonatal. *Rev Bras Enferm*; 2011 nov-dez 64(6):997-1002.
51. Freitas, EMDE; Nunes, ZB. O enfermeiro na prática de cateter central de inserção periférica em neonato / The nurse in the practice of peripherally inserted central catheter in neonatal care. *REME rev. min. enferm*; 2009 abr-jun 13(2): 215-224.

Received on: 13/05/2014

Reviews required: No

Approved on: 10/02/2015

Published on: 01/10/2016

Mailing address:

Cândida Caniçali Primo
Universidade Federal do Espírito Santo
Departamento de Enfermagem
Av. Marechal Campos, 1468, Maruípe, Vitória (ES), Brazil
ZIP Code: 29040-090